



GCSE
MATHEMATICS
8300/1H

Higher Tier Paper 1 Non-Calculator

Mark scheme
November 2023

Version: 1.0 Final

GRADEUP.UK



2 3 B G 8 3 0 0 / 1 H / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

GRADEUP.UK

Copyright information

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2023 AQA and its licensors. All rights reserved.

Glossary for Mark Schemes

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

M	Method marks are awarded for a correct method which could lead to a correct answer.
A	Accuracy marks are awarded when following on from a correct method. It is not necessary to always see the method. This can be implied.
B	Marks awarded independent of method.
ft	Follow through marks. Marks awarded for correct working following a mistake in an earlier step.
SC	Special case. Marks awarded for a common misinterpretation which has some mathematical worth.
M dep	A method mark dependent on a previous method mark being awarded.
B dep	A mark that can only be awarded if a previous independent mark has been awarded.
oe	Or equivalent. Accept answers that are equivalent. eg accept 0.5 as well as $\frac{1}{2}$
[a, b]	Accept values between a and b inclusive.
[a, b)	Accept values $a \leq \text{value} < b$
3.14 ...	Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416
Use of brackets	It is not necessary to see the bracketed work to award the marks.

Examiners should consistently apply the following principles.

Diagrams

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

Responses which appear to come from incorrect methods

Whenever there is doubt as to whether a student has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the student. In cases where there is no doubt that the answer has come from incorrect working then the student should be penalised.

Questions which ask students to show working

Instructions on marking will be given but usually marks are not awarded to students who show no working.

Questions which do not ask students to show working

As a general principle, a correct response is awarded full marks.

Misread or miscopy

Students often copy values from a question incorrectly. If the examiner thinks that the student has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

Further work

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

Choice

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

Work not replaced

Erased or crossed out work that is still legible should be marked.

Work replaced

Erased or crossed out work that has been replaced is not awarded marks.

Premature approximation

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.

Continental notation

Accept a comma used instead of a decimal point (for example, in measurements or currency), provided that it is clear to the examiner that the student intended it to be a decimal point.

Q	Answer	Mark	Comments
1	100	B1	

Q	Answer	Mark	Comments
2	60	B1	

Q	Answer	Mark	Comments
3	1, -2	B1	

Q	Answer	Mark	Comments
4	$a + 3$ or $3 + a$	B1	

GRADEUP.UK

Q	Answer	Mark	Comment	
5	$(8^2 \times 8 =) 8^3$ or $(8^9 \div 8^5 =) 8^4$ or 512 or 4096 or $8^2 \times 8 \div 8^9 \times 8^5$	M1		
	$(8^3$ or 512) \div (8^4 or 4096) or $8^{(2+1-9+5)}$ or $8^8 \times 8^{-9}$ or 8^{-1} or $\frac{1}{8}$	M1dep	oe in the form $8^n \div 8^{(n+1)}$ oe where index sums to -1 oe in the form $8^n \times 8^{(-n-1)}$ oe fraction	
	(0).125	A1		
	Additional Guidance			
	(0).125 and either 8^{-1} or $\frac{1}{8}$ on the answer line			M1M1A1
	(0).125 in working and 8^{-1} on the answer line			M1M1A0
	If a student attempts numerical and index working award the higher mark			

Q	Answer	Mark	Comment
	Valid description	B1	eg as downloads increase, so do CD sales downloads are about $\left[1\frac{1}{3}, 2\right]$ times as many as CDs CDs are about $\left[\frac{1}{2}, \frac{3}{4}\right]$ as many as downloads
Additional Guidance			
6a	Ignore 'Positive correlation'		
	Condone references to causality eg an increase in downloads causes an increase in CDs sold		B1
	As one goes up the other goes up / Both go up at a similar rate		B1
	They both go up		B0
	Downloads are always more than CDs		B0
	They are in direct proportion		B0
	The relationship is linear		B0

Q	Answer	Mark	Comment
6b	Alternative method 1 – reading from the graph		
	2.5(0) × 9000 or 22 500 or [5300, 5500]	M1	oe 2.5(0) may be 2 or 3 [5300, 5500] may be 5000
	2.5(0) × 9000 + 3 × [5300, 5500] or 22 500 + [15 900, 16 500]	M1dep	oe 2.5(0) may be 2 or 3 [5300, 5500] may be 5000
	[38 400, 39 000]	A1ft	ft 2 or 3 for 2.5(0) and/or 5000 for [5300, 5500]
	Alternative method 2 – using a multiplier		
	2.5(0) × 9000 or 22 500 or 9000 × [0.5, 0.75]	M1	oe 2.5(0) may be 2 or 3
	2.5(0) × 9000 + 3 × 9000 × [0.5, 0.75]	M1dep	oe 2.5(0) may be 2 or 3
	[36 000, 42 750] with 9000 × [0.5, 0.75] seen	A1ft	ft 2 or 3 for 2.5(0)
	Additional Guidance		
	Check graph for working		
	Working may be in pence, units not required for up to M2 Final answer in pence must have units to award A1		

Q	Answer	Mark	Comment	
7	Correct method to find 1%, 2%, 5%, 10%, 100% or 840% of the number	M1		
	Fully correct method	M1dep		
	600	A1		
	Additional Guidance			
	Up to M2 may be awarded for multiple attempts if no answer chosen			

GRADEUP.UK

Q	Answer	Mark	Comments
8	Alternative method 1: using algebra		
	$\frac{1}{2} \times k \times 12$ or $6k$ or $2 \times k + 3 \times m$ or $2k + 3m$	M1	oe correct expression for either area eg $2(k - 3) + 3(m + 2)$ or $k(m + 2) - m(k - 3)$
	$\frac{1}{2} \times k \times 12 = 2 \times k + 3 \times m$ or $4k = 3m$	M1dep	oe equation
	3 : 4	A1	oe ratio
	Alternative method 2: substituting a value for k		
	Substitutes a value for k and gives correct method or value for area of triangle or correct area in m for L-shape	M1	eg $k = 5$ and area of triangle is 30 or $k = 5$ and area of L-shape is $10 + 3m$
	Correct equation connecting the two areas or correct value of m for their k	M1dep	eg $k = 5$ and $30 = 10 + 3m$ or $k = 6$ and $m = 8$
	3 : 4	A1	oe ratio

Q	Answer	Mark	Comment	
9	$(x =) [2.25, 2.75]$ and $(x =) [9.25, 9.75]$	B2	B1 $(x =) [2.25, 2.75]$ or $(x =) [9.25, 9.75]$ or one or both values identified but not given in correct notation eg $(2.5, 0)$ and/or $(9.5, 0)$ or $2.5 < x < 9.5$ or 2.5 and/or 9.5 written on the graph or in working	
	Additional Guidance			
	$x =$ can be $x \approx$			
	[2.25, 2.75] and/or [9.25, 9.75] with one extra value		B1	
	[2.25, 2.75] and/or [9.25, 9.75] with more than one extra value		B0	
	Answer from use of formula or completing the square		B0	

Q	Answer	Mark	Comment	
10	$(\pi \times) \left(\frac{\sqrt{17}}{2}\right)^2$	M1	oe condone missing brackets	
	$\frac{17}{4}(\pi)$ or $4\frac{1}{4}(\pi)$ or $4.25(\pi)$	A1	oe fraction, mixed number or decimal	
	$(\pi \times) 5^2$ or $(\pi \times) 25$ or $\frac{60}{360}$ used	M1	oe	
	$\frac{25}{6}(\pi)$ or $4\frac{1}{6}(\pi)$ or $4.1(6\dots)(\pi)$ or $4.17(\pi)$	A1	oe fraction, mixed number or decimal	
	A with values in comparable form or A by $\frac{1}{12}(\pi)$ or A by $0.08(3\dots)(\pi)$	A1	eg values $\frac{51}{12}(\pi)$ and $\frac{50}{12}(\pi)$ $4\frac{1}{4}(\pi)$ and $4\frac{1}{6}(\pi)$ $4.2(5)(\pi)$ and $4.1(6\dots)(\pi)$ $4.2(5)(\pi)$ and $4.17(\pi)$ accept 'circle' for A allow comparison of fraction or decimal parts only if integer parts shown as equal	
	Additional Guidance			
	For the final mark, presence or absence of π must be the same for both values			
Accept consistent use of a numerical value of π throughout. The value can be 3 or 3.1 or 3.14 or 3.142 or better				

Q	Answer	Mark	Comment
11	$(x + 6)(x - 4)$	B2	either order B1 $(x + a)(x + b)$ where $ab = -24$ or $a + b = 2$
	Additional Guidance		
	Condone a multiplication sign between the brackets		
	Condone missing final bracket		
	Ignore an attempt to solve $(x + 6)(x - 4) = 0$		

Q	Answer	Mark	Comment
12(a)	2000	B1	

Q	Answer	Mark	Comment
12(b)	0.5 or $\frac{2 \times 10^3}{5 \times 10^{-1}}$ or $\frac{\text{their 2000}}{5 \times 10^{-1}}$ or $0.4 \times 10^{3 - (-1)}$ or 0.4×10^4	M1	oe their 2000 from part (a)
	4000 or 4×10^3	A1ft	ft $2 \times$ their 2000 in part (a)

Q	Answer	Mark	Comments
13	$10x + 5d$ or $10x = cx$ or $5d = 30$	M1	
	$10x = cx$ and $5d = 30$ or $c = 10$ or $d = 6$	M1dep	
	$c = 10$ $d = 6$	A1	

Q	Answer	Mark	Comments
14	$7x = 2x + 20$ or $\frac{x-20}{x} = \frac{2}{7}$ or $\frac{x+20}{x} = \frac{7}{2}$ or $20 \div (7 - 2)$ or $20 \div 5$ or 4	M1	oe any letter
	$7 \times 20 \div (7 - 2)$ or 28 or $2 \times 20 \div (7 - 2)$ or 8 or $(7 + 2) \times 20 \div (7 - 2)$	M1dep	oe hours for History hours for French total hours
	36	A1	
Q	Answer	Mark	Comments
15(a)	16	B1	

Q	Answer	Mark	Comments
15(b)	Average		
	Indicates that the statement is correct and uses the median to support their decision	B2	eg Yes, as median is lower (in 2019) B1 median is lower in 2019 with no or incorrect decision or median (for 2019) is 22.5
	Consistency		
	Indicates that the statement is correct and uses the IQR to support their decision or states that both quartiles were 1 minute greater in 2020	B2	eg IQRs are equal, so same consistency B1 IQR is the same with no or incorrect decision or IQR (for 2019) is 6 or IQR (for 2020) is 6
	Additional Guidance		
	Average		
	For B2 or B1, ignore reference to the LQ or UQ		
	Condone 22.30 for the median		
	For B2 or B1, if using 'average' for 'median' the value(s) must be given		
	Incorrect value for the 2019 median can score B1 if lower than 24 eg It's correct, the 2019 median was 23.5		B1
	Reference to IQR		B1max
	Median is 22.5, so true		B2
	Correct, the median is 1.5 lower in 2019 / Yes, it was 1.5 seconds faster		B2
	Consistency		
	Allow calculation seen to imply IQR eg (2019) $26 - 20 = 6$		
	Ignore reference to the range in 2019		
	Reference to median		B1max
	$26 - 20 = 27 - 21$ so this is true		B2
6 not accompanied by a calculation or reference to IQR		B0	
6 stated to be the range		B0	

Q	Answer	Mark	Comments
16(a)	$2x + x + 15 + 8 = 80$ or $3x + 23 = 80$ or $80 - 15 - 8$ or 57 or $\frac{x+8}{80}$ or $\frac{x+8}{3x+23}$	M1	oe implied by $\frac{57}{80}$
	$(x =) \frac{80-15-8}{3}$ or $\frac{57}{3}$ or 19 or 27	M1dep	oe implied by $\frac{19}{80}$
	$\frac{27}{80}$ or 0.3375 or 33.75%	A1	oe fraction, decimal or percentage
	Additional Guidance		
	Ignore incorrect conversion after correct answer seen		

Q	Answer	Mark	Comments
16(b)	$\frac{8}{23}$ or 0.3478... or 34.78...% or 0.348 or 34.8%	B1	oe fraction, decimal or percentage
	Additional Guidance		
	Ignore incorrect conversion after correct answer seen		

Q	Answer	Mark	Comments	
17	$2 \leq x \leq 5$ or $6 \leq 3x \leq 15$ or $x = 5$ or $3x = 15$	M1	may be in two parts implied by ($y =$) 27 or ($x =$) 2, 3, 4, 5 or ($3x =$) 6, 9, 12, 15	
	$y - 3 \times 5 < 12$ or $y < 12 + 3 \times 5$ or $y < 27$ or $y - 3 \times 5 \leq 11$ or $y \leq 11 + 3 \times 5$ or $y \leq 26$	M1dep	oe may be seen in a double-sided inequality eg condone $18 < y < 27$ using \leq or $=$	
	26	A1	SC1 17	
	Additional Guidance			
	SC1 is for the use of 2 instead of 5			
	All inequalities may be reversed, eg $2 \leq x \leq 5$ may be $5 \geq x \geq 2$			

Q	Answer	Mark	Comments	
18(a)	Rotation about (1, 0)			
	90° anticlockwise or 270° clockwise	B1	oe description of rotation condone 90° counter-clockwise eg quarter turn anticlockwise	
	Translation			
	$\begin{pmatrix} -2 \\ -6 \end{pmatrix}$	B1	oe description of translation eg 2 left and 6 down condone missing brackets SC1 B0B0 and point (-1, 4) identified	
	Additional Guidance			
	Condone missing degrees sign			
	(-2, -6)			B0
	Compound transformation			B0 for that part

Q	Answer	Mark	Comments
18(b)	(4, 4) and (7, 7) or (1, 1) and (6, 6)	B1	condone (5, 5) and (10, 10) either order

19	Alternative method 1		
	(0.4̇ =) $\frac{4}{9}$ or $10x - x = 4$ or $9x = 4$	M1	oe eg $100x - x = 44$ or $99x = 44$ or $100x - 10x = 40$ or $90x = 40$ any letter
	$\frac{7}{9} \div 10$ or $\frac{7}{90}$ or $10y - y = 0.7$ or $9y = 0.7$	M1	oe eg $100y - y = 7.7$ or $99y = 7.7$ or $100y - 10y = 7$ or $90y = 7$ any letter
	$\frac{47}{90}$	A1	oe single fraction
	Alternative method 2		
	0.52̇	M1	oe
	$10x - x = 4.7$ or $9x = 4.7$	M1dep	oe eg $100x - x = 51.7$ or $99x = 51.7$ or $100x - 10x = 47$ or $90x = 47$ any letter
	$\frac{47}{90}$	A1	oe single fraction
	Additional Guidance		
	For M marks, allow fractions with decimal numerator or denominator eg in alt 1, $\frac{0.7}{9}$ scores M1 and in alt 2, $\frac{4.7}{9}$ scores M2		

Q	Answer	Mark	Comments
20	(x =) 60 or (3x =) 180 or (y =) 45 or (2y =) 90	M1	implied by sin 60 or tan 45 must be selected and not just in a table
	3 × 60 – 2 × 45 or 180 – 90 or (w =) 90 or cos 90	M1dep	
	0 with M2 awarded	A1	
	Additional Guidance		
	0 with no working		M0M0A0
	Condone degrees sign on answer with correct working		M1M1A1

Q	Answer	Mark	Comments	
21	$2(4x)^2 + 9$	M1	oe	
	$32x^2 + 9$	M1dep	dep on 2nd M1 may be implied by 4th mark	
	$8x + 9$	M1	may be implied by 4th mark	
	$32x^2 - 8x = 0$ or $32x^2 = 8x$	M1	oe equation with brackets expanded rearranges their $f^{-1}(x) =$ their $gh(x)$ to correctly collect terms	
	0 and $\frac{1}{4}$	A1	oe eg 0 and $\frac{8}{32}$	
	Additional Guidance			
	With no terms to collect in their equation the 4th mark cannot be awarded			
	$8x + 9 = 8x^2 + 36$ $0 = 8x^2 - 8x + 27$		MOMOM1 M1A0	
	$8x - 9 = 8x^2 + 36$ $0 = 8x^2 - 8x + 45$		MOMOM0 M1A0	
	$8x + 9 = 2(4x)^2 + 9$ $8x + 9 = 16x^2 + 9$ $8x = 16x^2$		M1MOM1 M1A0	
$8x + 9 = 4x(2x^2 + 9)$ $0 = 8x^3 + 28x - 9$		MOMOM1 M1A0		

Q	Answer	Mark	Comments
22	Alternative method 1		
	$\frac{180}{9+3}$ or 15 or $3 \times \frac{180}{9+3}$ or 45 or $9 \times \frac{180}{9+3}$ or 135	M1	oe eg $180 \times \frac{1}{3+1}$ or $180 \div 4$ size of angle c size of angle a
	$5 \times \frac{180}{9+3}$ or or 5×15 or 75 or $7 \times \frac{180}{9+3}$ or $(3 + 5 + 9) \times \frac{180}{9+3}$ or 255	M1dep	oe size of angle b
	105	A1	
	Alternative method 2		
	$a : b : c : d = 9 : 5 : 3 : 7$ or $b : d = 5 : 7$	M1	oe implied by $d = 7$
	$\frac{7}{9+5+3+7} \times 360$ or $\frac{7}{24} \times 360$ or $\frac{7}{5+7} \times 180$ or $\frac{7}{12} \times 180$ or 7×15	M1dep	oe allow numerator to be 5
	105	A1	

Q	Answer	Mark	Comments
23	Alternative method 1		
	$\frac{7\sqrt{3}}{\sqrt{20}} \times \frac{\sqrt{20}}{\sqrt{20}}$	M1	oe eg $\frac{7\sqrt{3}}{\sqrt{2}\sqrt{10}} \times \frac{\sqrt{2}\sqrt{10}}{\sqrt{2}\sqrt{10}}$
	$\frac{7\sqrt{60}}{20}$	M1dep	oe single rationalised fraction
	$\frac{7\sqrt{15}}{10}$ or $\frac{14\sqrt{15}}{20}$	A1	oe in the form $\frac{x\sqrt{15}}{y}$ where x and y are integers
	Alternative method 2		
	$\frac{7\sqrt{3}}{2\sqrt{5}}$	M1	
	$\frac{7\sqrt{3}}{2\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}}$	M1dep	oe rationalisation eg $\frac{7\sqrt{3}}{2\sqrt{5}} \times \frac{2\sqrt{5}}{2\sqrt{5}}$
	$\frac{7\sqrt{15}}{10}$ or $\frac{14\sqrt{15}}{20}$	A1	oe in the form $\frac{x\sqrt{15}}{y}$ where x and y are integers

Mark scheme and Additional Guidance continue on the next page

Q	Answer	Mark	Comments
23 cont	Alternative method 3		
	$\frac{7}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}}$ or $\frac{7\sqrt{2}}{2}$ or $\frac{\sqrt{3}}{\sqrt{10}} \times \frac{\sqrt{10}}{\sqrt{10}}$ or $\frac{\sqrt{30}}{10}$	M1	oe
	$\frac{7\sqrt{2}}{2} \times \frac{\sqrt{30}}{10}$ or $\frac{7\sqrt{60}}{20}$	M1dep	oe rationalised
	$\frac{7\sqrt{15}}{10}$ or $\frac{14\sqrt{15}}{20}$	A1	oe in the form $\frac{x\sqrt{15}}{y}$ where x and y are integers
	Alternative method 4		
	$\frac{7}{\sqrt{2}} \times \frac{\sqrt{3}}{\sqrt{10}} \times \frac{\sqrt{5}}{\sqrt{5}}$	M1	oe
	$\frac{7\sqrt{5}}{\sqrt{10}} \times \frac{\sqrt{3}}{\sqrt{10}}$ or $\frac{7}{\sqrt{2}} \times \frac{\sqrt{15}}{\sqrt{50}}$ or $\frac{7\sqrt{15}}{\sqrt{100}}$	M1dep	oe one term or product of two terms with numerator $7\sqrt{15}$
	$\frac{7\sqrt{15}}{10}$ or $\frac{14\sqrt{15}}{20}$	A1	oe in the form $\frac{x\sqrt{15}}{y}$ where x and y are integers

Q	Answer	Mark	Comments
24	$\frac{1}{2}$	B1	oe

Q	Answer	Mark	Comments
25	$r^2 = \frac{8}{9}$ or $\sqrt{\frac{8}{9}}$ or $\frac{2\sqrt{2}}{\sqrt{9}}$ or $\frac{\sqrt{8}}{3}$ or $\frac{2\sqrt{2}}{3}$ or $\left(\sqrt{\frac{8}{9}}\right)^3$ or $\frac{8\sqrt{8}}{27}$	M1	oe eg $\left(\frac{8}{9}\right)^{\frac{1}{2}}$ allow $\pm\sqrt{\frac{8}{9}}$ etc
	$\frac{16\sqrt{2}}{27}$	A1	oe in the form $\frac{c\sqrt{2}}{d}$ where c and d are integers

Q	Answer	Mark	Comments
26(a)	$\sqrt[4]{5\frac{1}{16}}$ or $\left(5\frac{1}{16}\right)^{\frac{1}{4}}$ or $\frac{81^{\frac{1}{4}}}{16^{\frac{1}{4}}}$ or $\left(81^{\frac{1}{4}} = \right) 3$ or $\left(16^{\frac{1}{4}} = \right) 2$	M1	oe eg $\sqrt[4]{\frac{81}{16}}$ condone missing brackets
	$\frac{3}{2}$ or $1\frac{1}{2}$ or 1.5	A1	

Q	Answer	Mark	Comments
26(b)	$(7^{2m})^{2.5}$ or $49^{2.5m}$ or $(7^2)^{2.5m}$	M1	oe with base changed to 7 and/or brackets removed
	7^{5m}	A1	

Q	Answer	Mark	Comments
27	$-4 < x < 4$ or $4 > x > -4$	B1	condone the inequality in two parts eg $x > -4$ and $x < 4$

Q	Answer	Mark	Comments	
28	$(\overrightarrow{XR}) = \frac{3}{2}(2\mathbf{a} + 4\mathbf{b})$ or $3\mathbf{a} + 6\mathbf{b}$ or $(\overrightarrow{QR}) = \frac{5}{2}(2\mathbf{a} + 4\mathbf{b})$ or $5\mathbf{a} + 10\mathbf{b}$	M1	oe may be on diagram	
	$(\overrightarrow{PS}) = -5\mathbf{a} + 2\mathbf{a} + 4\mathbf{b}$ $+ \frac{3}{2}(2\mathbf{a} + 4\mathbf{b}) + \mathbf{a} - 8\mathbf{b}$ or $(\overrightarrow{PS}) = -5\mathbf{a} + \frac{5}{2}(2\mathbf{a} + 4\mathbf{b}) + \mathbf{a} - 8\mathbf{b}$ or $(\overrightarrow{PS}) = \mathbf{a} + 2\mathbf{b}$	M1dep	oe may be on diagram	
	$(\overrightarrow{PS}) = \mathbf{a} + 2\mathbf{b}$ and indication why PS is parallel to QR	A1	eg $2(\mathbf{a} + 2\mathbf{b}) = 2\mathbf{a} + 4\mathbf{b}$ or $5\mathbf{a} + 10\mathbf{b} = 5(\mathbf{a} + 2\mathbf{b})$ or $\mathbf{a} + 2\mathbf{b}$ and \overrightarrow{QR} is a multiple of \overrightarrow{PS}	
	Additional Guidance			
	Some or all vectors may be reversed and the final mark can be from using a negative constant eg $\overrightarrow{RX} = -3\mathbf{a} - 6\mathbf{b}$ $\overrightarrow{PS} = \mathbf{a} + 2\mathbf{b}$ $-3(\mathbf{a} + 2\mathbf{b}) = -3\mathbf{a} - 6\mathbf{b}$			M1 M1 A1

Q	Answer	Mark	Comments
29(a)	$y = \cos x$ passes through $(180^\circ, -1)$	B1	

Q	Answer	Mark	Comments
29(b)	None of the graphs pass through $(270^\circ, 1)$	B1	