



---

# GCSE MATHEMATICS 8300/3F

Foundation Tier Paper 3 Calculator

---

Mark scheme

June 2022

---

Version: 1.0 Final

GRADEUP.UK



2 2 6 G 8 3 0 0 / 3 F / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [aqa.org.uk](http://aqa.org.uk)

GRADEUP.UK

#### **Copyright information**

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2022 AQA and its licensors. All rights reserved.

**Glossary for Mark Schemes**

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

<b>M</b>	Method marks are awarded for a correct method which could lead to a correct answer.
<b>A</b>	Accuracy marks are awarded when following on from a correct method. It is not necessary to always see the method. This can be implied.
<b>B</b>	Marks awarded independent of method.
<b>ft</b>	Follow through marks. Marks awarded for correct working following a mistake in an earlier step.
<b>SC</b>	Special case. Marks awarded for a common misinterpretation which has some mathematical worth.
<b>M dep</b>	A method mark dependent on a previous method mark being awarded.
<b>B dep</b>	A mark that can only be awarded if a previous independent mark has been awarded.
<b>oe</b>	Or equivalent. Accept answers that are equivalent. eg accept 0.5 as well as $\frac{1}{2}$
<b>[a, b]</b>	Accept values between a and b inclusive.
<b>[a, b)</b>	Accept values $a \leq \text{value} < b$
<b>3.14 ...</b>	Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416
<b>Use of brackets</b>	It is not necessary to see the bracketed work to award the marks.

Examiners should consistently apply the following principles.

**Diagrams**

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

**Responses which appear to come from incorrect methods**

Whenever there is doubt as to whether a student has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the student. In cases where there is no doubt that the answer has come from incorrect working then the student should be penalised.

**Questions which ask students to show working**

Instructions on marking will be given but usually marks are not awarded to students who show no working.

**Questions which do not ask students to show working**

As a general principle, a correct response is awarded full marks.

**Misread or miscopy**

Students often copy values from a question incorrectly. If the examiner thinks that the student has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

**Further work**

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

**Choice**

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

**Work not replaced**

Erased or crossed out work that is still legible should be marked.

**Work replaced**

Erased or crossed out work that has been replaced is not awarded marks.

**Premature approximation**

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.

**Continental notation**

Accept a comma used instead of a decimal point (for example, in measurements or currency), provided that it is clear to the examiner that the student intended it to be a decimal point.

Q	Answer	Mark	Comments
1	25%	B1	

Q	Answer	Mark	Comments
2	5	B1	

Q	Answer	Mark	Comments
3	$\frac{9}{100}$	B1	

Q	Answer	Mark	Comments
4	8c	B1	

Q	Answer	Mark	Comments
5(a)	centimetres or millimetres or inches	B1	allow abbreviations eg cm, mm, in
	grams or milligrams or ounces	B1	allow abbreviations eg g, mg, oz
	<b>Additional Guidance</b>		
	Mark intention eg condone ou or incorrect spellings		
	Ignore any numbers with correct units		

Q	Answer	Mark	Comments
5(b)	20 + 40 or 60 or 90 or 1 (h) + 1 (h) + 30 (m) or 150 or 2 (h) 30 (m)	M1	
	$2\frac{1}{2}$ or 2.5	A1	oe answer in hours eg two and a half SC1 2.3(0)
	<b>Additional Guidance</b>		
	Ignore rounding attempt to 2 or 3 after correct answer seen eg 2.5 in working with answer 2		M1A1
	2 h 30 min in working with answer 2		M1A0
	1.9(0)		M0

Q	Answer	Mark	Comments
6	$8 \times (0.)60$ or 480 or 4.8(0)	M1	oe
	10 – their 4.8(0) or 5.2(0) or 1000 – their 480 or 520	M1	oe $0.6(0) \leq \text{their } 4.8(0) < 10$ $60 \leq \text{their } 480 < 1000$ 5.2(0) or 520 implies M2
	26	A1	
	<b>Additional Guidance</b>		
	Up to M2 may be awarded for correct work, with no or incorrect answer, even if this is seen amongst multiple attempts		
	$60 \div 8 = 7.50$ then $10 - 7.50$		M0M1A0

Q	Answer	Mark	Comments
7(a)	3	B1	allow answer in words

Q	Answer	Mark	Comments
7(b)	<b>Alternative method 1</b>		
	2 + 6 + 9 or 17 (2008) or 5 + 8 + 3 or 16 (2012)	M1	oe
	17 and 16	A1	
	<b>Alternative method 2</b>		
	2 – 5 + 6 – 8 + 9 – 3 or –3 – 2 + 6 or 5 – 2 + 8 – 6 + 3 – 9 or 3 + 2 – 6	M1	oe eg 3 more gold, 2 more silver, 6 fewer bronze
	Indication that there was 1 more medal in 2008	A1	oe indication there was 1 less in 2012
	<b>Additional Guidance</b>		
	17 must not be linked with 2012, 16 must not be linked with 2008		
	Ignore further work after correct answer seen		

Q	Answer	Mark	Comments
<b>7(c)</b>	Valid reason	B1	eg 25 is not a multiple of 3 or 25 ÷ 3 is not a whole number or 8 + 8 + 8 = 24 or 9 + 9 + 9 = 27
	<b>Additional Guidance</b>		
	Ignore incorrect or irrelevant statements alongside correct statements, unless contradictory		
	3 is not a factor of 25		B1
	(25 ÷ 3 =) 8.3(...)		B1
	(25 ÷ 3 =) $8\frac{1}{3}$		B1
	3 × 8 = 24 or 3 × 9 = 27		B1
	It would have to be 8, 8 and 9		B1
	25 divided by 3 is a decimal		B1
	25 can't be (fully) divided by 3 (condone)		B1
	3 doesn't go into 25 (condone)		B1
	25 doesn't fit evenly into 3 (condone)		B1
	The three equal totals would not add up to 25		B1
	None of the equal totals would add up to 25		B0
	There are not 3 whole numbers that add to make 25		B0
25 is not a factor of 3		B0	
The difference between the possible answers is 3		B0	

Q	Answer	Mark	Comments
<b>8</b>	<b>Alternative method 1</b>		
	5.6 ÷ 7 or 0.8	M1	oe
	5.6 + their 0.8 or 6.4	M1	oe their 0.8 must not be 0.4 and must be less than 5.6
	their 6.4 – 6 or 0.4	M1dep	oe dep on 2nd M1
	400	A1	SC1 any correct conversion litres to millilitres with M0 scored
	<b>Alternative method 2</b>		
	5.6 × 1000 or 5600 or 6 × 1000 or 6000 or 5.6 ÷ 7 or 0.8	M1	oe
	their 5600 ÷ 7 or their 0.8 × 1000 or 800	M1	oe their 5600 must include the digits 56 consecutively their 0.8 must not be 0.4 and must be less than 5.6
	their 5600 + their 5600 ÷ 7 or their 5600 + their 0.8 × 1000 or 6400	M1dep	oe their 5600 must include the digits 56 consecutively their 0.8 must not be 0.4 and must be less than 5.6 dep on 2nd M1
	400	A1	SC1 any correct conversion litres to millilitres with M0 scored

**Additional Guidance continues on the next page**

		<b>Additional Guidance</b>	
<b>8 cont</b>		Up to M3 may be awarded for correct work, with no or incorrect answer, even if this is seen amongst multiple attempts	
		Beware of 0.4 or 400 from incorrect working	
		6400 or 0.4 (not from incorrect working)	M1M1M1
		0.9 and 6.5 and 0.5 or 0.9 and 6.5 and 500 (500 implies 0.5)	M0M1M1A0
		$560 \div 7$ and $560 + 80$ (560 includes the digits 56 consecutively)	M0M1M1A0
		560 and 80 and 640	M0M1M1A0
		560 and 600 and 80 and 40	M0M1M1A0
		In Alt 2, $0.0056 \div 7$ (0.0056 includes the digits 56 consecutively)	M0M1

GRADEUP.UK

Q	Answer	Mark	Comments
9(a)	8 in Time exercising Less than 1 hour	B1	
	23 in Exercise taken No	B1	
	58 in Total number of students	B1ft	ft 35 + their 23 or 27 + their 8 + their 23
	<b>Additional Guidance</b>		
	8 in Time exercising Less than 1 hour		B1
	47 in Exercise taken No		B0
	82 in Total number of students		B1ft
	7 in Time exercising Less than 1 hour		B0
	25 in Exercise taken No		B0
	59 in Total number of students		B1ft

Q	Answer	Mark	Comments
9(b)	$\frac{27}{35}$ or 0.77(...) or 77(. ...)%	B1	oe fraction
	<b>Additional Guidance</b>		
	Ignore attempts to simplify or convert after correct fraction seen		
	eg1 $\frac{27}{35}$ seen, answer $\frac{5}{7}$		B1
	eg2 $\frac{27}{35}$ seen, answer 7.7%		B1
	Ignore words if correct answer seen		
	eg1 $\frac{27}{35}$ seen, answer 27 out of 35		B1
	eg2 77%, unlikely		B1
Answer given as ratio (even if correct answer also seen)			
eg 27 : 35		B0	
Answer only in words			
eg 27 out of 35		B0	
Only 77 (without %)		B0	

Q	Answer	Mark	Comments
10(a)	Hexagon	B1	

Q	Answer	Mark	Comments
10(b)	Valid reason	B1	eg sides are not equal or angles are not equal
	<b>Additional Guidance</b>		
	Ignore incorrect or irrelevant statements alongside correct statements, unless contradictory		
	There are no lines of symmetry		B1
	It has reflex angles		B1
	Regular polygons must have equal sides		B1
	All sides are different (condone)		B1
	Some sides are more than 1 cm		B1
	It doesn't have a line of symmetry		B1
	It doesn't have one line of symmetry		B0

Q	Answer	Mark	Comments
10(c)	2	B1	allow in words
	4	B1	allow in words

Q	Answer	Mark	Comments
11(a)	4	B1	
	<b>Additional Guidance</b>		
	4 in output oval with answer line blank		B1
	4 in output oval with different answer on answer line		B0

Q	Answer	Mark	Comments
11(b)	$d = 3c - 5$ or $d = 3 \times c - 5$	B2	oe eg $d = -5 + 3c$ B1 $d = 3c \dots$ or $d = 3 \times c \dots$ or $3c - 5$ or $3 \times c - 5$ SC1 $c = \frac{d+5}{3}$
	<b>Additional Guidance</b>		
	Further incorrect work after a B2 response is B1 eg $d = 3c - 5$ followed by $d = -15c$		B1
	Further incorrect work after a B1 response is B1 eg $3c - 5$ followed by $-15c$		B1
	Condone $3c - 5$ on answer line if $d = 3c - 5$ seen in working		B2
	$3c - 5 = d$		B2
	$d = c \times 3 - 5$		B2
	$d = c3 - 5$		B1
$c3 - 5$		B0	

Q	Answer	Mark	Comments
12(a)	$3x + 2y$	B2	either order B1 $3x$ or $2y$
	<b>Additional Guidance</b>		
	Further incorrect work after a B2 response is B1 eg $3x + 2y$ followed by $5xy$		B1
	Further incorrect work after a B1 response is B1 eg $15x + 2y$ followed by $30xy$		B1

Q	Answer	Mark	Comments
12(b)	$8 \times 25$ or 200 or $25^2$ or 625	M1	oe
	$8 \times 25$ or 200 and $25^2 - b$ or $625 - b$ or $25^2 - 8 \times 25$ or $625 - 200$	M1dep	oe may be seen in an equation
	425	A1	
	<b>Additional Guidance</b>		
	Embedded answer		M1M1A0

Q	Answer	Mark	Comments
12(c)	$3w + 5$	B1	

Q	Answer	Mark	Comments
13	True Cannot tell	B2	B1 one correct
	<b>Additional Guidance</b>		
	A tick and a cross in the same row – mark the tick		
	Allow any unambiguous indication		

Q	Answer	Mark	Comments
14(a)	8	B1	

Q	Answer	Mark	Comments
14(b)	$1 \times 7$ and $2 \times 5$ and $3 \times 4$ and $4 \times 1$ and $5 \times 3$ or $7$ and $10$ and $12$ and $4$ and $15$ or $48$	M1	oe allow one error or omission
	$(7 + 10 + 12 + 4 + 15) \div 20$ or $48 \div 20$ or their $48 \div 20$	M1dep	oe eg $\frac{48}{20}$ or $\frac{12}{5}$ or $2\frac{2}{5}$ without working their 48 must be the correct sum of their products
	2.4	A1	SC1 33.75
	<b>Additional Guidance</b>		
	$48 \div 5$		M1M0
	$1 \times 7 + 2 \times 5 + 3 \times 4 + 4 \times 1 + 5 \times 5$ (5 × 5 is one error) $58 \div 20 = 2.9$		M1 M1A0
	$8 + 10 + 12 + 4 + 15$ (8 is one error) $49 \div 20 = 2.45$		M1 M1A0
	Answer 2 after 2.4 seen		M1M1A0
	$7 + 10 + 12 + 4 + 15 \div 20$ not recovered		M1M0
	Correct products or values seen but a different method used is a choice of methods eg 7 10 12 4 15 followed by $20 \div 5$ or $20 \div 15$		M0

Q	Answer	Mark	Comments
15(a)	300 or 360 or 480 or 7 ( $\times$ 60) or 7th or any 3 multiples of 60 that are greater than 60	M1	
	420	A1	
	<b>Additional Guidance</b>		
	420 in working with answer 7 or 7th or $7 \times 60$		M1A0

Q	Answer	Mark	Comments	
15(b)	6	B2	B1 answer 2 or answer 3 or answer 2 ( $\times$ ) 3 or answer 2, 6 or answer 3, 6 or answer 2, 3, 6 or (1) 2 3 4 6 (12) or (1) 2 3 6 9 (18) or (12 $\Rightarrow$ ) 2 ( $\times$ ) 2 ( $\times$ ) 3 or $2^2$ ( $\times$ ) 3 or (18 $\Rightarrow$ ) 2 ( $\times$ ) 3 ( $\times$ ) 3 or 2 ( $\times$ ) $3^2$	
			<b>Additional Guidance</b>	
			If correct answer 6 is obtained from a list of factors, then the list must contain no errors	
			For use of prime factors, allow in repeated division or a factor tree or a Venn diagram or inclusion of 1	
			List of factors may be seen in factor pairs (allow repeats) eg (1 $\times$ 12) 2 $\times$ 6 3 $\times$ 4	B1

Q	Answer	Mark	Comments
16	$2 \times 3.5$ or 7	M1	oe implied by 5.7(...) or 5 r5 or 42
	Ticks No and 5.7(...) or Ticks No and 42	A1	oe eg $\frac{40}{7}$ is less than 6
	<b>Additional Guidance</b>		
	Ignore area and volume calculations		
	Ticks No and 5 r5		M1A1
	Ticks No and $5\frac{5}{7}$		M1A1
	Ticks No and 2 cm too short		M1A1
	Ticks Yes and 5.7(...)		M1A0
$12 \times 3.5$		M1	

Q	Answer	Mark	Comments
17(a)	3200	B1	

Q	Answer	Mark	Comments
17(b)	12	B1	

Q	Answer	Mark	Comments
18	$b$ and $c$	B1	

Q	Answer	Mark	Comments		
19(a)	Straight line from (0, 0) to (10, 35)	B2	$\pm \frac{1}{2}$ square		
			B1 one correct point $\pm \frac{1}{2}$ square		
			from (2, 7) to (10, 35) seen or plotted or		
			one correct ratio apart from 2 : 7 or		
			one correct pair of amounts apart from 2 juice 7 water		
			<b>Additional Guidance</b>		
			Mark intention		
			If no points plotted, a correct point from (2, 7) to (10, 35) can be implied by a straight line with positive gradient		
	Two points plotted with the same $x$ -coordinate is choice unless the line is drawn through one of the points				
Condone straight line from (2, 7) to (10, 35)	B2				
(2, 7) seen with graph not drawn or incorrect	B1				
10 : 35 seen with graph not drawn or incorrect	B1				
6 juice 21 water with graph not drawn or incorrect	B1				

Q	Answer	Mark	Comments
19(b)	<b>Alternative method 1 – uses the given ratio</b>		
	17.5	B1	
	<b>Alternative method 2 – uses their graph</b>		
	Correct water reading for 5 litres of juice from their straight line	B1ft	$\pm \frac{1}{2}$ square
	<b>Additional Guidance</b>		
	17 or 18 from a correct straight line		B1

Q	Answer	Mark	Comments
<b>20</b>	Ticks Yes and valid reason	B1	eg ticks Yes and she has thrown more often
	<b>Additional Guidance</b>		
	Ignore incorrect or irrelevant statements alongside correct statements, unless contradictory		
	Ticks No	B0	
	Ticks Yes and 60 is more than 40	B1	
	Ticks Yes and 60 is 20 more than 40	B1	
	Ticks Yes and 60 is 10 more than 40	B1	
	(ignore incorrect value 10)		
	Ticks Yes and she has more data to look at	B1	
	Ticks Yes and her number of throws is higher	B1	
	Ticks Yes and Bianca used more throws which gives her a higher chance of getting heads	B1	
	Ticks Yes and Adam has less number of throws and has more heads (ignore irrelevant has more heads)	B1	
	Ticks Yes and Bianca throws more coins	B1	
	Ticks Yes and she threw it 60 times, Adam only 40	B1	
	Ticks Yes and she threw it 60 times, Adam 40	B0	
	Ticks Yes and she threw it 60 times and got 20	B0	
	Ticks Yes and the probability is $\frac{20}{60}$	B0	
Ticks Yes and because her total is higher	B0		

Q	Answer	Mark	Comments
21	<b>Alternative method 1</b>		
	tan identified	M1	oe eg $\tan^{-1}$
	$\tan x = \frac{10}{4}$ or $\tan x = \frac{5}{2}$ or $\tan x = 2.5$	M1dep	oe eg $\tan^{-1} \frac{10}{4}$ or $90 - \tan^{-1} \frac{4}{10}$
	[68, 68.2]	A1	SC1 [21.8, 22]
	<b>Alternative method 2</b>		
	$\sin x = \frac{10}{\sqrt{4^2 + 10^2}}$ or $\cos x = \frac{4}{\sqrt{4^2 + 10^2}}$	M2	oe eg $\sin x = \frac{10}{\sqrt{116}}$ or $\sin^{-1} \frac{10}{\sqrt{4^2 + 10^2}}$ or $\cos x = \frac{4}{\sqrt{116}}$ or $\cos^{-1} \frac{4}{\sqrt{4^2 + 10^2}}$ or $90 - \sin^{-1} \frac{4}{\sqrt{4^2 + 10^2}}$ or $90 - \cos^{-1} \frac{10}{\sqrt{4^2 + 10^2}}$
	[68, 68.2]	A1	SC1 [21.8, 22]
	<b>Additional Guidance</b>		
	Accept 10.77 or 10.8 or $2\sqrt{29}$ for $\sqrt{116}$		
	Tan can be identified by, for example, circling TOA in SOHCAHTOA		
Answer from accurate drawing		M0M0A0	
$\sin x = \frac{10 \sin 90}{\sqrt{116}}$		M2	
$(x =) \tan 2.5$ or $(x =) \tan 0.4$ or $(x =) \tan \left(\frac{10}{4}\right)^{-1}$ unless recovered		M1M0A0	
$\tan = \frac{10}{4}$ or $\tan = \frac{4}{10}$ or $\tan x = \frac{4}{10}$ with no further correct working		M1M0A0	

Q	Answer	Mark	Comments
	3 + 2 or 5 and $5\frac{1}{2} + 3\frac{1}{2}$ or 9 or $5\frac{1}{2} - 3$ or $2\frac{1}{2}$ and $3\frac{1}{2} - 2$ or $1\frac{1}{2}$ or 4	M1	oe eg 180 + 120 or 300 and 330 + 210 or 540 implied by $5\frac{1}{2} + 3\frac{1}{2} - 3 - 2$
22	$\frac{9-5}{5}$ or $\frac{2\frac{1}{2}+1\frac{1}{2}}{3+2}$ or $\frac{4}{5}$ or 0.8 or $\frac{5\frac{1}{2}+3\frac{1}{2}}{3+2} (\times 100)$ or $\frac{9}{5} (\times 100)$ or 1.8 ( $\times 100$ ) or 180	M1dep	oe eg $\frac{5\frac{1}{2}+3\frac{1}{2}-3-2}{3+2}$ eg $\frac{540-300}{300}$ or $\frac{240}{300}$ or 1.8 – 1
	80	A1	
<b>Additional Guidance</b>			
Allow working fully in minutes but units must be consistent in a single calculation eg 2 h 30 and 1 h 30 eg 3 + 2 = 5 and 330 + 210 = 540 eg 3 + 120 and $330 + 3\frac{1}{2}$ unless recovered			M1 M1 M0
$3 + 2 = 6$ , $5\frac{1}{2} + 3\frac{1}{2} = 9$ , $9 - 6 = 3$ , 3 = 50%			M1M1A0
$3 + 2 = 6$ , $5\frac{1}{2} + 3\frac{1}{2} = 9$ , answer 50% (3 is implied)			M1M1A0
$9 - 6 = 3$ , 3 = 50% (no method shown for 6)			M0M0A0

Q	Answer	Mark	Comments
23(a)	-1 and 5	B1	either order
	<b>Additional Guidance</b>		
	Ignore $x =$ written before answers		
	(-1, 0) or (5, 0)		B0

Q	Answer	Mark	Comments
23(b)	(2, -9)	B2	B1 $x = 2$ or (2, ...) or $y = -9$ or (... , -9) or $(x - 2)^2 - 9$
			B1ft correct $y$ -coordinate for their $x$ -coordinate with $x \neq -1, 0$ or 5 SC1 (-9, 2)
	<b>Additional Guidance</b>		
	If answer line is blank, check diagram for indication of $x$ or $y$ values		
	(3, -9)		B1
	(3, -8)		B1ft
	(1, -8)		B1ft
	(2.5, -8.75)		B1ft
(0, -5)		B0ft	

Q	Answer	Mark	Comments	
24	(8th term $\Rightarrow$ ) $2^8$ or 256	M1	oe may be implied	
	Common difference of A indicated as 3	M1	may be implied eg $3n \dots$ or $\dots + 3(n - 1)$	
	$3n + 10 =$ their 256 or (their 256 $-$ 10) $\div$ 3 or (their 256 $-$ 13) $\div$ 3 or 81	M1dep	oe equation eg $13 + 3(n - 1) = 2^8$ dep on 2nd M1 their 256 may be any number and may be in index form	
	82	A1		
	<b>Additional Guidance</b>			
	$n + 3$ implies 2nd M1			
	Do not award M1 for 256 if it is in a list of powers of 2 unless it is indicated or it is the highest power evaluated			
	Common difference of 3 may be shown on the progression for the 2nd M1			
	10, (13, 16, 19, 22), 25 without common difference of 3 shown does not imply 2nd M1			
	82 from trial and improvement		M3A1	
	Embedded answer $3 \times 82 + 10 = 256$		M3A0	
	$3n + 10 = 256$ or $3n + 10 = 2^8$ or $3n = 246$		M1M1M1	
	$3n - 10 = 256$		M1M1M0	
	$3n + 10 = 16$ ( $2^8$ not seen)		M0M1M1	
	$3n + 6 = 2^8$		M1M1M0	
$256 - 22 = 234$ , $234 \div 3$ (indicating common difference of 3)		M1M1M0		
$3n - 8 = 128$ ( $2^8$ not seen)		M0M1M0		

Q	Answer	Mark	Comments
25	<b>Alternative method 1</b>		
	$4 \times 2$ or 8	M1	oe may be seen in an equation eg $3 \times x + 4 \times 2 = 44$
	$\frac{44 - 4 \times 2}{3}$ or $\frac{36}{3}$ or 12	M1dep	oe
	38	A1	
	<b>Alternative method 2</b>		
	$7 \times 2$ or 14	M1	oe may be seen in an equation eg $7 \times 2 + 3 \times y = 44$
	$\frac{44 - 7 \times 2}{3}$ or $\frac{30}{3}$ or 10	M1dep	oe
	38	A1	
	<b>Additional Guidance</b>		
	Up to M2 may be awarded for correct work, with no or incorrect answer, even if this is seen amongst multiple attempts		
	Working for up to M2 may be seen on the diagram		
	Beware of 38 from incorrect working $7 + 3 + 7 + 3 = 20$ , $7 + 2 + 7 + 2 = 18$ , $20 + 18 = 38$		MOMOA0

Q	Answer	Mark	Comments
26	$\begin{pmatrix} 5 \\ 23 \end{pmatrix}$	B1	
	<b>Additional Guidance</b>		
	Condone $\begin{pmatrix} 5 \\ 23 \end{pmatrix}$		B1

Q	Answer	Mark	Comments
27	$330 \div (3 + 2)$ or $330 \div 5$ or 66	M1	oe eg $\frac{330}{5}$
	their $66 \times 2$ or 132	M1dep	oe $\frac{2}{5} \times 330$ scores M2
	$294 \div 7$ or 42 or $294 \div 7 \times 3$ or 126	M1	oe eg $\frac{294}{7}$ or $\frac{3}{7} \times 294$
	132 and 126 and A	A1	
	<b>Additional Guidance</b>		
	132 and 88.2 and A		M1M1M0A0

Q	Answer	Mark	Comments
28	<b>Alternative method 1 – compares speeds in m/s</b>		
	$200 \div 24$ or $8.3(3\dots)$	M1	oe eg $\frac{200}{24}$ or $8\frac{1}{3}$
	$28.8 \times 1000 \div 60 \div 60$ or 8	M1	oe eg $28800 \div 3600$ or $28.8 \div 3.6$
	8 and $8.3(3\dots)$ and Tom	A1	oe eg 8 and $8\frac{1}{3}$ and Tom
	<b>Alternative method 2 – compares speeds in km/h</b>		
	$200 \div 24$ or $8.3(3\dots)$	M1	oe eg $\frac{200}{24}$ or $8\frac{1}{3}$
	their $8.3(3\dots) \div 1000 \times 60 \times 60$ or 30	M1dep	oe eg $0.0083(3\dots) \times 3600$
	30 and Tom	A1	
	<b>Alternative method 3 – time for Adil starting with m/s</b>		
	$28.8 \times 1000 \div 60 \div 60$ or 8	M1	oe eg $28800 \div 3600$
	$200 \div$ their 8 or 25	M1dep	oe eg $\frac{200}{8}$
	25 and Tom	A1	oe eg Tom by 1s
	<b>Alternative method 4 – time for Adil starting with km/h</b>		
	$\frac{200 \div 1000}{28.8}$ or $[0.0069, 0.007]$ or $\frac{200}{28.8}$ or $[6.9, 7]$	M1	oe eg $\frac{0.2}{28.8}$  eg $\frac{125}{18}$
	their $[0.0069, 0.007] \times 60 \times 60$ or their $[6.9, 7] \div 1000 \times 60 \times 60$ or 25	M1dep	oe eg $\frac{0.2}{28.8} \times 3600$
	25 and Tom	A1	oe eg Tom by 1s

Mark scheme and Additional Guidance continue on the next page

Q	Answer	Mark	Comments
28 cont	<b>Alternative method 5 – distance for Adil in 24s</b>		
	28 800 × 24 or 691 200 or 28.8 ÷ 60 ÷ 60 or 0.008 or 28.8 × 24 or 691.2	M1	oe eg $\frac{3456}{5}$
	their 691 200 ÷ 60 ÷ 60 or their 0.008 × 1000 × 24 or their 691.2 × 1000 ÷ 60 ÷ 60 or 192	M1dep	oe eg 28 800 × 24 ÷ 3600
	192 and Tom	A1	
	<b>Additional Guidance</b>		
	Up to M2 may be awarded for correct work, with no or incorrect answer, even if this is seen amongst multiple attempts		
	Ignore all units		
	Allow other correct comparisons eg 500 and 480 (this is metres per minute) eg 500 and 480 and Tom		M1M1 M1M1A1
	200 m = 0.2 km, 24 s = 24 ÷ 60 ÷ 60 = $\frac{1}{150}$ hour, 0.2 ÷ $\frac{1}{150}$ = 30 and Tom		M1M1A1
	$\frac{200 \div 1000}{24} = \frac{1}{120}$ (or 0.0083...)		M1

Q	Answer	Mark	Comments
29	3.55 ≤ mass < 3.65	B1	